

STRAWBERRY PLANT NAMED 'CARMELA'

BOTANICAL CLASSIFICATION

The present invention relates to a new *Fragaria x ananassa* Duch. plant.

VARIETAL DENOMINATION

5 The new plant has the varietal denomination 'CARMELA'.

BACKGROUND OF THE INVENTION

The new variety of strawberry was created in a breeding program by crossing two parents; in particular, by crossing as seed parent an undistributed strawberry parent designated 86-032 (~~unpatented~~ not patented) and as pollen parent an undistributed strawberry
10 parent designated 9261 (~~unpatented~~ not patented). Female is a component of a parent collection, from a selection made between plants issued from seeds in a free pollination in a population of different origin. Male is a selection from breeder's program of Planasa. Both parental varieties are proprietary and have not been commercialized, distributed or patented.

The resulting seedling of the new variety was grown and asexually propagated by
15 runners in Soria, Spain, 3°W., 41°N., 3000 feet elevation. Clones of the new variety were further asexually propagated and extensively tested. This propagation and testing has demonstrated that the combination of traits disclosed herein which characterize the new variety are fixed and retained true to type through successive generations of asexual reproduction.

20 SUMMARY OF THE INVENTION

The present invention relates to a new and distinct strawberry variety. The varietal denomination of the new variety is 'CARMELA'. Among the characteristics which appear to distinguish the new variety from other varieties are a combination of traits which include inflorescence that appears level with the foliage and abundant production of orange red
25 colored, conical shaped, and very firm fruit, medium fruit size, and very early time of ripening (50% of plants with ripe fruits).

COMPARISON TO CLOSEST VARIETY

The new variety is closest to the variety 'Tudnew' (U.S. Plant Pat. No. 10,960), but is distinguished therefrom by the following characteristics possessed by 'Carmela' which are different than, or not possessed by, 'Tudnew' (U.S. Plant Pat. No. 10,960).

5 1. 'Tudnew' (U.S. Plant Pat. No. 10,960) exhibits a habit of plant flat globose than 'Carmela' exhibits a habit of plant globose.

 2. 'Tudnew' (U.S. Plant Pat. No. 10,960) shows a leaf color of upper side (RHS green group near 138B to 138A) less dark than 'Carmela' (RHS green group near 131B to 135A).

10 3. Length/width ratio of the terminal leaflet in 'Tudnew' (U.S. Plant Pat. No. 10,960) is as long as broad, than in 'Carmela' it is longer than broad.

 4. In 'Tudnew' (U.S. Plant Pat. No. 10,960) the position of the inflorescence relative to foliage is above . In 'Carmela' it is level with.

 5. The attitude of hairs in 'Tudnew' (U.S. Plant Pat. No. 10,960) is slightly
15 outwards, than in 'Carmela' it is upwards.

 6. The fruit size of 'Tudnew' (U.S. Plant Pat. No. 10,960) is more large than of 'Carmela'.

 7. 'Tudnew' (U.S. Plant Pat. No. 10,960) shows a red fruit color (RHS red group near 42C to 42A), whereas in 'Carmela' it is an orange red fruit color (RHS orange red group
20 near 33A to 34A).

 8. The predominant shape of fruits of 'Tudnew' (U.S. Plant Pat. No. 10,960) is biconical, whereas in 'Carmela' it is conical.

 The differences in the leaf color of upper side of the new variety and 'Tudnew' (U.S. Plant Pat. No. 10,960) are shown in FIGS. 2 and 3. The differences in the fruits of the new
25 variety and 'Tudnew' (U.S. Plant Pat. No. 10,960) are shown in FIGS. 7, 8, and 9,10. These differences are maintained during the harvest season.

 9. 'Tudnew' (U.S. Plant Pat. No. 10,960) shows a firmness fruit slightly more than 'Carmela' fruit.

 10. Precocity in 'Tudnew' (U.S. Plant Pat. No. 10,960) is smaller than in 'Carmela'.

30 BRIEF DESCRIPTION OF ILLUSTRATIONS

The accompanying photographs show typical specimens of the new variety, designated 98.08V.134 in the illustrations, including fruit, foliage and flower, in color as nearly true as it is reasonably possible to make in color illustrations of this character.

5 The plants depicted in the drawings were planted in October in the farm of La Mogalla in Cartaya (Huelva), Spain, about 7°W, 37°N, 45 feet elevation.

Drawings were taken in April: minimum temperate about 16°C to 18°C, maximum temperate about 31 to 33° C.

10 FIG. 1 shows the undersurface of a typical terminal leaflet of the new variety (designated 98.08V.134) with a shape of base obtuse and in which the length/width ratio is longer than broad.

FIG. 2 shows the top and undersurface of a complete leave of the new variety (designated 98.08V.134). FIG 3 shows the top of a complete leave of 'Tudnew' (U.S. Plant Pat. No. 10,960) (designated 92.H1.51). In them we can see that the leaf color of upper side of 'Tudnew' (U.S. Plant Pat. No. 10,960) is less dark than 'Carmela'.

15 FIG. 4 and 5 show the fruit against a background of the top surface of the foliage of the new variety (designated 98.08V.134).

FIG. 6 shows the flower and reproductive organs of the new variety (designated 98.08V.134).

20 FIG. 7 and 8 show typical fruit of the new variety (designated 98.08V.134) whole, sliced and in cross section, illustrating the typical flesh and flesh coloration, conspicuous core and core cavity, conical shape and orange red fruit color.

FIG. 9 and 10 show typical fruit of the 'Tudnew' (U.S. Plant Pat. No. 10,960) (designated 92.H1.51) whole, sliced and in cross-section, illustrating the typical flesh and flesh coloration, conspicuous core and core cavity, biconical shape and red fruit color.

DESCRIPTION OF THE NEW VARIETY

The following detailed description of the new variety is based upon observations taken of plants and fruits grown “under glass”, *i.e.*, under tunnel, in the farm of La Mogalla in Cartaya (Huelva), Spain, 7°W., 37°N., 45 feet elevation.

- 5 The following description is in accordance with UPOV terminology and the color terminology herein is in accordance with *The Royal Horticultural Society Colour Chart* (R.H.S.C.C.). The color descriptions and other phenotypical descriptions may deviate from the stated values and descriptions depending upon variation in environmental, seasonal, climatic and cultural conditions.

10 PROPAGATION

The new variety is principally propagated by way of runners. Although propagation by runners is presently preferred, other know methods of propagating strawberry plants may be used. Strawberries root well after transplanting.

- 15 The term “blistering” used herein refers to the texture or rugosity or surface undulation inherent to leaves and is generally a constant characteristic.

GENERAL

- 'Carmela' is a short day variety that needs an induction to flowering by chilling, such as occurs at a high elevation nursery (fresh plant) or with cold storage (referred to as a frigo). Usually a short time is sufficient. 'Carmela' is self-fertile. It produces large quantity of
20 pollen throughout the seasons and pollination is generally good as there are very few malformed fruit.

Production: Plants described are from high elevation nursery.

- 25 Description taken in Cartaya (Huelva), Spain of plants planted in October 2002.

 Number of repetitions: 2

 Plants per repetition: 225

- Comparison with 'Tudnew' (U.S. Plant Pat. No. 10,960): The new variety is compared
30 with “Tudnew” (U.S. Plant Pat. No.10,960) (designated 92.H1.51) in FIG. 2 and FIG. 3 and FIG. 7 and FIG. 8 and FIG. 9 and FIG. 10.

Accumulated production of 1st quality fruit (g/plant)

	Variety	21-Fb	28-Mr	25-Apr	15-May
5	MILSEI	12	98	451	802
	CAMAROSA	15	96	519	1035
10	TUDNEW	23	102	481	936
	CARMELA	48	168	458	809

	Variety	1st + 2nd Quality Fruit	Total	Weight (g/fruit)
15	MILSEI	802 + 109	911	22 - 21
20	CAMAROSA	1035 + 183	1218	24 - 23
	TUDNEW	936 + 104	1040	24 - 23
25	CARMELA	809 + 90	899	22 - 21

Production total, to 15 May, of First Quality Fruit (1st quality)
and Second Quality Fruit (2nd quality) in g/plant

	Variety	1 st quality	2 nd quality	TOTAL (1 st quality + 2 nd quality)	% 2 nd quality
30	MILSEI	802	109	911	12
35	CAMAROSA	1035	183	1218	15
	TUDNEW	936	104	1040	10
40	CARMELA	809	90	899	10

$$\% 2^{\text{nd}} \text{ quality} = \frac{2^{\text{nd}} \text{ quality}}{\text{TOTAL}} \times 100$$

Weight (g/Fruit) at two dates: 28 March and 15 May

	WEIGHT (g/fruit)	28 March	15 May
5	MILSEI	22	21
	CAMAROSA	24	23
	TUDNEW	24	23
10	CARMELA	22	21

WEIGHT is shown as the average weight per fruit in First Quality Fruits.

FRUIT ANALYSIS

		CAMAROSA	CARMELA (98.08V.134)	MILSEI	TUDNEW
20	Firmness (KG)	0.60	0.86	0.34	0.98
25	Humidity & Volatile Matter (%)	91.10	90.90	92.00	91.80
	Dry Matter (%)	8.90	9.10	8.00	8.20
30	PH (to20°)	3.50	3.60	3.60	3.60
	Acidity as Anhydride Citric (%)	1.10	1.10	1.00	1.00
35	Soluble Solids (°Brix)	7.90	8.10	7.20	6.70
	Maturity Index	7.30	7.70	7.10	6.80
40	Content in Ascorbic Acid (ppm)	670	600	560	600
	Dominant Tonality(nm)	495	495	500	500

Luminosity:

Transmittance to 460nm	11.10	38.60	20.90	41.30
------------------------	-------	-------	-------	-------

The following definitions apply:

5

Firmness: It is the fruit's resistance to penetration measured in Kilograms (Kg). The measure given has been obtained by the penetrometer ROZE Mod. Arbelette, with a 50 mm² section head.

10 Dry Matter: It is the weight of the residual left from the trituration of the fruit after the drying process at a temperature of 103°C ± 2°C until reaching constant weight.

$$(\%) \text{ Dry Matter} = \frac{\text{Weight Dry Matter}}{\text{Weight Fresh Matter}} \times 100$$

15

Humidity & Volatile Matter: Represents the content in volatile matters and water of the fruits.

$$(\%) \text{ Humidity \& Volatile Matter} = 100 - \% \text{ Dry Matter}$$

20

Maturity Index: Relation between Soluble solids and Acidity as Anhydride Citric.

$$\text{Maturity Index} = \frac{\text{Soluble solids}}{\text{Acidity as Anhydride Citric}}$$

25

The following additional information is provided to further describe the new variety.

Variety: Carmela. Breeder Ref. 98.08V.134

Classification: *Fragaria x ananassa* Duch.

30

Plant:

Habit.- Globose.

Density.- Medium.

Vigor.- Medium.

35

Height.- About 21 cm.

Width.- About 22 cm.

Leaf:

Upper side.- RHS green group color (near 131B to 135 A)

Underside.- RHS green group color (near 143D to 138D).

5 Length.- About 10 cm.

Width- section.- ~~Slightly concave~~ About 13 cm. to 14 cm.

Leaf surface undulation or Blistering.- Very strong.

Number of leaflets.- Three only.

Leaf stem characteristics:

10 Color.- RHS green group (near 138D).

Position of hairs.-upwards.

Length.- About 12 cm.

Terminal leaflet:

Length/width ratio.- Longer than broad.

15 Length.- About 7,5 cm.

Width.- About 6 cm.

Shape of base.-Obtuse.

Shape of teeth.- Crenate.

Petiole:

20 Position of hairs.- Upwards.

Length.- About 12 cm.

Stipule:

Anthocyanin coloration.- Strong. Color.- RHS greyed-red group coloration (near 178C to 179A)

25 Stolons:

Number.- Many, about 9 to 10.

Thickness.- Medium, about 3.2 mm.

Pubescence.- Medium.

Color.- RHS green group (near 138D to 142D).

Inflorescence:

Position relative to foliage.- Level with.

Flower:

Size.- Medium.

5 Size of calyx relative to corolla.- Larger.

Spacing of petals.- Overlapping.

Flower characteristics:

Diameter primary flowers.- About 2.0 – 2.5 cm.

Diameter secondary flowers.- About 2.0 cm.

10 Number of petals.- Normally about 6. No significant fragrance.

Time from bloom to mature fruit (in Huelva, Spain).- About 33 to 37 days.

Stamens.- Numerous with pollen present, fertile and abundant.

Length – approximately 4 mm. Color – RHS white group (near 155D to 155C).

15 Anthers.- Generally average in size. Color – RHS yellow group (near 12B to 13B) and darkening with advanced maturity.

Pollen.- Fertile and abundant. Color – RHS yellow orange group (near 14B to 15B)

Pistils.- Numerous, generally average in size. Color – RHS yellow group (near 12B to 13B).

20 Petal:

Length/width ratio.- As long as broad.

Fruiting truss:

Attitude.- Semi-erect.

25 Fruit:

Ratio of length/maximum width.- Much longer than broad.

Color.- RHS orange red group (near 33A to 34A).

Peduncle length of inflorescence stem.- Primary fruit about 9 to 11 cm, secondary fruit about 6 to 7 cm, color near 138D.

Primary fruit:

Length.- About 5.0-5.5 cm.

Width.- About 3.0-3.5 cm.

Secondary fruit:

5 Length.- About 4.5-5.0 cm.

Width.- About 2.5-3.0 cm.

Size.- Medium.

Predominant shape.- Conical.

Difference in shapes between primary and secondary fruits.- Slight.

10 Band without achenes.- Medium.

Color of achenes.- RHS orange red group (near 33C to 33B).

Unevenness of surface.- Absent or very weak.

Evenness of color.- Even.

Glossiness.- Medium.

15 Insertion of achenes.- Below surface.

Insertion of calyx.- Set above fruit.

Pose of the calyx segments.- Reflexed.

Size of calyx in relation to fruit diameter.- Much larger.

Adherence of calyx.- Very strong.

20 Firmness.- Firm.

Color of flesh.- RHS orange red group (near 34C to 34B), lightening toward center.

Distribution of red color of flesh.- Marginal and central.

Hollow center.- Weakly expressed.

Sweetness.- Strong.

25 Acidity.- Medium.

Time of flowering (50% of plants at first flower).- Very early.

Time of ripening (50% of plants with ripe fruits)- Very early.

Type of bearing.- Not remontant.

Chilling.- Weak.

Planting date.- October 30, 2002.

10% Flowering.- December 23, 2002.

First mature fruits.- January 26, 2003

Maturity (15-20 gms/plant).- February 8, 2003

- 5 Time of flowering data: Date of planting: October 30, 2002 in the farm of La Mogalla, in Cartaya (Huelva), Spain, about 7°W, 37°N, 45 feet elevation. 10% flowering occurs about December 23, 2002 with first mature fruit about January 26, 2003 and maturity (15-20 g / plant) about February 8, 2003.

- 10 Time of flowers (50% of plants at first flower): about December 30, 2002

Storage Qualities: 'Carmela' fruit maintain their quality characteristics when keeping them in a frigo chamber at temperatures of about 2°C. during 48 hours. The fruit's color remains substantially the same.

- 15 Time of ripening: After planting as aforesaid, plants are grown in raised beds under tunnel (small tunnel with small holes in plastic walls). Water and fertilizer were applied through drip irrigation. Time of ripening (50% of plants with ripe fruit) is about January 31, 2003. First mature fruit is about January 26, 2003 and maturity (15-20 gms/plant) is about
20 February 8, 2003.

- General: The growing period in Huelva, Spain, where the observations were made, is between about December,10 and May,15 of each year, with a maximum production at about mid-April. 'Carmela' is a short variety that benefits from induction to flowering by chilling,
25 usually a few hours are sufficient, preferably at temperatures of 7°C. or less. Normally, the minimum number of hours is accumulated in the field during several days.

Disease resistance: No particular sensitivity to any disease or parasite has been observed for 'Carmela'.

I CLAIM:

1. A new and distinct strawberry plant of the variety substantially as shown and described.

ABSTRACT

Described is a strawberry variety having inflorescence about level with the foliage and abundant fruit production with early ripening.

5

10

TXT IRV1077458.1-* -06/24/04 4:13 PM